

DOI: <https://doi.org/10.47391/JPMA.748>

Dynamic changes of coagulation and fibrinolytic biomarkers in perioperative arthroplasty patients

Cong Wang¹, Songjie Ji², Zhifang Chen³, Zhiwei Liu⁴, Heng Zhou⁵, Chunmin Li⁶, Bin Zhao⁷, Chunsheng Li⁸

1,3,4,5,7 Department of Emergency, Beijing Jishuitan Hospital, Peking, China;

2,6 Department of Orthopaedic, Beijing Jishuitan Hospital, Peking, China; **8** Department of Emergency, Beijing Chao-Yang Hospital, Capital Medical University, Peking, China

Correspondence: Chunsheng Li **Email:** cslemu@163.com

Abstract

Objective: To evaluate Coagulation and fibrinolytic parameters after total joint arthroplasty (TJA) and provide evidence for optimization of timing of perioperative anticoagulation medicine.

Methods: The prospective study was conducted at the Jishuitan Hospital of Peking University from January to April in 2016, and comprised patients who were scheduled consecutively to undergo primary total knee arthroplasty (TKA) or total hip arthroplasty (THA). Blood samples were obtained at day 1 preoperatively and day1, day 3 postoperatively. Antigenic levels of protein C (PC), endothelial protein C receptor (EPCR), tissue factor pathway inhibitor (TFPI), antithrombin III (AT-III), plasminogen activator inhibitor 1 (PAI-1) and

tissue plasminogen activator (tPA) were measured with commercially available enzyme-linked immunosorbent assay kits.

Results: Postoperative levels of coagulation parameters TFPI and AT-III were increased compared to preoperative values (118.7 ± 34.6 vs 70.0 ± 20.5 $\mu\text{g/ml}$ for AT-III, and 26.37 ± 7.91 vs 16.68 ± 8.92 $\mu\text{g/l}$ for TFPI), while postoperative levels of coagulation parameters PC and EPCR were decreased (0.88 ± 0.30 vs 2.03 ± 0.66 $\mu\text{g/ml}$ for PC, and 100.8 ± 31.0 vs 199.4 ± 57.4 $\mu\text{g/ml}$ for EPCR). Postoperative levels of fibrinolytic parameter tPA was increased compared to preoperative values (2.87 ± 0.83 vs 2.03 ± 1.03 $\mu\text{g/l}$), while its specific inhibitor PAI-1 was decreased (0.88 ± 0.30 vs 2.03 ± 0.66 $\mu\text{g/l}$).

Conclusion: These results demonstrated the perturbation of the coagulation and fibrinolytic system of patients undergoing TJA. Hypercoagulation and hyperfibrinolysis were observed in postoperative patients, which suggested anticoagulant therapy is effective and necessary.

Keywords: Total joint arthroplasty, anticoagulation, fibrinolysis.

Introduction

Total joint arthroplasty (TJA), including total knee arthroplasty (TKA) and total hip arthroplasty (THA), is an effective treatment in patients with severe joint diseases. It was well determined that patients with TJA had a high risk of postoperative bleeding events and venous thromboembolism (VTE)¹⁻³. TJA is associated with a risk of thromboembolism because patients undergoing this

46 procedure had a hypercoagulable state frequently. According to previous studies,
 47 the incidence of VTE in patients with TJA could be as high as 15~30 %, which
 48 led to a mortality rate of 2 % because of threatening pulmonary embolism ¹⁻³.
 49 It was also shown that perioperational thrombotic events induced a
 50 hypercoagulable state and increased fibrinolytic activity after TKA ^{4, 5}.
 51 Understanding of the coagulable and fibrinolytic dysregulation will be
 52 important to have more therapies to prevent thrombotic events in TJA patients.
 53 In 2013, the new guideline recommended that anticoagulant therapy should be
 54 used for at least 14 days after TJA to prevent VTE ². The use of anticoagulant
 55 agents (such as warfarin and low molecular weight heparin) protected patients
 56 from potential thrombotic events. The mechanisms for postoperative
 57 coagulation dysregulation are not fully understood. Protein C (PC) is a major
 58 component of the human anticoagulation system, which regulates thrombosis
 59 and limits inflammatory response ⁶. Protein C circulates in the blood in the form
 60 of inactive zymogen. During surgical stress, PC on the surface of endothelial
 61 cells is activated by binding thrombin and thrombomodulin ⁷. Endothelial cell
 62 protein C receptor (EPCR) can augment the
 63 thrombin-thrombomodulin-mediated PC activation ⁸. (AT-III), the inhibitor for
 64 serine proteases, is the most important physiological anticoagulant in the plasma
 65 ⁹. Tissue factor pathway inhibitor (TFPI), also known as an exogenous pathway
 66 inhibitor (EPA) and lipoprotein-related coagulation inhibitor (LACI), is a major
 67 inhibitor during the extrinsic coagulation pathway ¹⁰. TFPI is an endogenous

68 anticoagulant protein produced primarily by endothelial cells and secondarily
 69 from liver and monocytes/macrophages ¹⁰. It was shown that TFPI can be
 70 combined with heparin sulfate on the surface of endothelial cells. Heparin
 71 injection could release TFPI from endothelial cells, so it was considered that the
 72 increased TFPI in plasma was correlated with the routine application of
 73 low-molecular-weight heparin for preventive anticoagulation ^{10, 11}.
 74 complications, including allergic reactions, infectious agent contamination,
 75 transfusion related acute lung injury and hemolytic complications ¹²⁻¹⁴.
 76 Increased fibrinolytic activity in blood samples after TKA might be attributed to
 77 increased fibrin degradation, which leads to excessive bleeding ¹⁵. Tissue
 78 plasminogen activator (tPA) is an important element of the fibrinolytic system ¹⁶.
 79 tPA has a high affinity towards the fibrin-plasminogen complex within the
 80 thrombus, which enables it to rapidly form into fibrin-tPA-plasminogen
 81 complex. This process greatly facilitates the activation of plasminogen and
 82 exerts a highly efficient and specific thrombolytic effect ¹⁷. The production and
 83 release of tPA was decreased in the injured vascular endothelial cells, which
 84 induce thrombosis and decreased fibrinolytic activity ¹⁸. Plasminogen activator
 85 inhibitor 1 (PAI-1) is a specific inhibitor of tPA, which is produced and released
 86 primarily by vascular endothelial cells and partially by platelets ^{19, 20}.
 87 All of the above basic biomarkers are used for further clinical application.
 88 Throughout the coagulation process, a small quantity of activated coagulation
 89 factors could activate a large number of cascaded downstream coagulation

factors, so the whole coagulation process shows a huge amplification phenomenon. Antithrombin-, protein C- or protein S- deficiency constitutes a major risk factor for venous thromboembolism ^{21, 22}. Understanding of the coagulable and fibrinolytic dysregulation would be important to have more therapies to prevent thrombotic events and bleeding complications in patients after TJA surgery. In this research, we evaluated the coagulation functions and fibrinolytic parameters during the perioperative period, which provide evidence for the time window of anticoagulation treatment after TKA surgery.

Patients and Methods

A total of 119 patients enrolled in the study were scheduled consecutively to undergo primary THA or TKA at the Jishuitan Hospital of Peking University from January to April in 2016. Only patients with surgical indications for THA or TKA participated in this study. TKA were performed in 48 patients and THA were performed in 71 patients. The surgical indications included instability, pain, deformity, and dysfunction of the hip or knee joint. Patients who meet any of the following criteria were considered not eligible for the study: (1) patients with cerebral or gastrointestinal bleeding within the previous 6 months; (2) ischemic heart disease in the previous 6 months; (3) hepatic failure; (4) end-stage kidney disease; (5) coagulation or fibrinolysis disorder; (6) any kind of cancer. Before their enrollment, all patients were provided a detailed explanation of the risks and alternatives to participation in this research and

informed consent was provided according to the Declaration of Helsinki. This study was approved by the institutional review board of Beijing Jishuitan Hospital.

Blood samples were obtained from peripheral veins on the preoperative day, postoperative 24h and postoperative 72h respectively. Samples were centrifuged at 3000 rpm for 15 minutes at room temperature. Plasma samples were stored at -80 °C. Protein levels of antithrombin III, protein C, endothelial protein C receptor, tissue factor pathway inhibitor, plasminogen activator inhibitor 1 and tissue plasminogen activator were measured from these plasma samples with enzyme linked immunosorbent assay (ELISA) kit (Elabscience, E-EL-H0432c for antithrombin III, E-EL-H1167c for protein C, E-EL-H0065c for endothelial protein C receptor, E-EL-H0165c for tissue factor pathway inhibitor, E-EL-H2104c for plasminogen activator inhibitor 1, E-EL-H2106c for tissue plasminogen activator). Incidences of high-risk pulmonary embolism and DVT were estimated through inquiring and vascular ultrasound with a 3-month follow-up.

Continuous data were expressed as mean \pm standard deviation and compared with Student's t-test or one-way analysis of variance (ANOVA). Categorical data were presented as percentages and frequencies and compared with chi-square test. All statistical analyses were performed with SPSS 19.0 software. P-values<0.05 were considered statistically significant.

Results

A total of 119 patients (30 male /89 female, mean age: 57.7 ± 12.01 years, mean body mass index: 24.3 ± 6.67 kg/m²) were enrolled in this study. The clinical characteristics of patients with TJA are shown in Table 1. Among the 119 patients, 42 suffered from hypertension, 8 patients were in presence of diabetes, 7 diagnosed with coronary heart disease, 2 diagnosed with cerebrovascular disease, 2 experienced joint replacement surgery and 3 were glucocorticoids users. The levels of PC and its receptor EPCR were measured in TJA patients before and 24h, 72h after the procedure, respectively (Table 2). PC was significantly decreased in postoperative 24h and 72h groups compared to preoperative group (Fig. 1A), while the change of the expression of EPCR is consistent with PC (Fig. 1B). The reduced expression levels of both PC and its receptor validated the hypercoagulatable state in TJA patients postoperatively. It was suggested that reduced bioactivity of the PC anticoagulation system was involved in the pathogenesis of thrombosis event in TJA patients. The level of another important physiological anticoagulant factor AT-III was also investigated (Table 2). An increasing level of AT-III expression was found at the postoperative state (Fig. 2A). The postoperative TFPI expression demonstrated an increasing trend when compared with the preoperative level, which indicated an antithrombosis tendency (Fig. 2B).

As a physiological activator for human fibrinolysis system that is synthesized by the endothelial cells, tPA could transfer plasminogen into plasmin and

dissolve thrombus. Compared to the preoperative state, the expression of PAI-1 was significantly reduced while that of tPA was increased (Table 2, Fig. 3), which indicated the state of hyperfibrinolysis following TJA.

Follow up at 3-month revealed that the morbidity of DVT was 5% (6/119), all ultrasound result were intravenous thrombus. No patient experienced high-risk pulmonary embolism. It illustrated that anticoagulation therapy is essential and effective.

Discussion

As mentioned previously, prevalence of DVT following TJA was high despite the wide use of anticoagulation measures. The factors of DVT formation included slow blood flow, vascular wall injury and hypercoagulatable state. Under a normal physiologic state, coagulation, hemostatic function and anticoagulation, fibrinolytic function of the body are in a dynamic equilibrium. Any alteration from either side would result in a pathological state or occurrence of disease. Strengthened coagulation, hemostatic function, reduced anticoagulation and fibrinolytic function would lead to thrombosis. Coagulation is induced by multiple procoagulant factors such as endothelial injury and tissue factor release, which is activated in a cascade manner by series of coagulation-related cytokines^{23, 24}. The studies of the pathogenesis of postoperative thrombotic events in TJA patients were of great importance for the prevention of thrombotic events after joint replacement. The changes of

178 coagulation and fibrinolysis system were observed in TJA patients before,
179 during and after the operation. Our research found that human body was in a
180 hypercoagulable and hyperfibrinolytic state after TJA. It was observed that the
181 expression of PC and its specific receptor EPCR were both decreased after the
182 operation, which implied the hypercoagulatable state in TJA patients
183 postoperatively. The plasma PAI-1 can promptly bind with tPA in the 1:1 ratio,
184 which then deactivates tPA ²⁵. Therefore, increase of PAI-1 can be observed at
185 the pre-thrombotic state and during thrombosis. The expression of tPA was
186 increased, while that of its inhibitor PAI-1 was decreased after the operation.
187 It was reported that surgical stress led to reduced regional release of AT-III
188 through impairing the endothelial cell function, which activated the intrinsic and
189 extrinsic coagulation pathways, consumed a lot of AT-III and eventually
190 weakened the anti-thrombotic function ²⁶. It has been shown that the extrinsic
191 coagulation pathway is mainly regulated by TFPI, a Kunitz-type serine protease
192 inhibitor ²⁷. Our results were opposite to the previous results. The following
193 reasons were considered. Direct anticoagulant effect of antithrombin was slow
194 and weak, but its anticoagulant effect can be enhanced for 2000 times after
195 combining with heparin. Physiologically, antithrombin enhanced the
196 anticoagulation of vascular endothelium by binding heparan sulfate on the
197 surface of endothelial cells. All of our patients received low-molecular-weight
198 heparin anticoagulation immediately after the operation. It might be due to the
199 use of anticoagulants, the expression of physiological anticoagulant substances

such as AT-III and TFPI were increased after surgery.

At present, conventional heparin, low-molecular-weight heparin, as well as vitamin K antagonists are proved to be effective for the prevention and treatment of thrombosis, which also have limitations²⁸. Currently, rivaroxaban, dabigatran and apixaban have been licensed to be used in patients after TJA operation²⁸. Current evidence suggested that rivaroxaban reduced the incidence of symptomatic VTE compared with enoxaparin, whereas drug-related bleeding events were also increased²⁹. Thus, new oral anticoagulants have become the focus of current research.

Conclusion

The reduced expression of PC and EPCR after TJA indicated PC anticoagulation system was inhibited and a hypercoagulatable state was present. Meanwhile, increased expression of a fibrinolytic system marker tPA and decreased expression of PAI-1 suggested a state of hyperfibrinolysis was observed following TJA. Multiple cytokines such as TFPI and tPA are all secreted by endothelial cells, which indicated that the changes may correlate with the injuries of vascular endothelial cells. Moreover, the investigation of new anticoagulants was also closely related to the pathophysiological process of postoperative coagulation, which was worthy of our further investigation and tracking.

222 **Disclaimer:** None to declare.

223 **Conflict of Interest:** None to declare.

224 **Funding Disclosure:** None to declare.

225

226 **References**

- 227 1. He ML, Xiao ZM, Lei M, Li TS, Wu H, Liao J. Continuous passive motion
228 for preventing venous thromboembolism after total knee arthroplasty. *Cochrane*
229 *Database Syst Rev* 2014; CD008207.
- 230 2. Lieberman JR, Pensak MJ. Prevention of venous thromboembolic disease
231 after total hip and knee arthroplasty. *J Bone Joint Surg Am* 2013; 95: 1801-1811.
- 232 3. Thorsberger M, Jensen T, Olesen J, Weeke P, Wahlsten L, Jensen P, et al.
233 Opportunities for improvement: anticoagulation in patients undergoing total
234 knee or hip replacement. *J Orthop Sci* 2015; 20: 1036-1045.
- 235 4. Guler N, Burleson A, Syed D, Banos A, Hopkinson W, Hoppensteadt D, et
236 al. Fibrinolytic Dysregulation in Total Joint Arthroplasty Patients: Potential
237 Clinical Implications. *Clin Appl Thromb Hemost* 2016; 22: 372-376.
- 238 5. Yukizawa Y, Inaba Y, Kobayashi N, Ike H, Kubota S, Saito T. Selective
239 pharmacological prophylaxis based on individual risk assessment using plasma
240 levels of soluble fibrin and plasminogen-activator inhibitor-1 following total hip
241 arthroplasty. *Mod Rheumatol* 2014; 24: 835-839.
- 242 6. Esmon NL, Safa O, Smirnov MD, Esmon CT. Antiphospholipid antibodies
243 and the protein C pathway. *J Autoimmun* 2000; 15: 221-225.

- 244 7. Esmon C. The protein C pathway. *Crit Care Med* 2000; 28: S44-48.
- 245 8. Clouse LH, Comp PC. The regulation of hemostasis: the protein C system.
246 *N Engl J Med* 1986; 314: 1298-1304.
- 247 9. Beresford CH, Owen MC. Antithrombin III. *Int J Biochem* 1990; 22:
248 121-128.
- 249 10. Bajaj MS, Bajaj SP. Tissue factor pathway inhibitor: potential therapeutic
250 applications. *Thromb Haemost* 1997; 78: 471-477.
- 251 11. Novotny WF. Tissue factor pathway inhibitor. *Semin Thromb Hemost* 1994;
252 20: 101-108.
- 253 12. Levine BR, Haughom B, Strong B, Hellman M, Frank RM. Blood
254 management strategies for total knee arthroplasty. *J Am Acad Orthop Surg* 2014;
255 22: 361-371.
- 256 13. Jaworski K, Maslanka K, Kosior DA. Transfusion-related acute lung injury:
257 a dangerous and underdiagnosed noncardiogenic pulmonary edema. *Cardiol J*
258 2013; 20: 337-344.
- 259 14. Dilek I, Demir C, Bay A, Akdeniz H, Oner AF. Seropositivity rates of
260 HBsAg, anti-HCV, anti-HIV and VDRL in blood donors in Eastern Turkey. *Turk*
261 *J Haematol* 2007; 24: 4-7.
- 262 15. Matsuda K, Nozawa M, Katsube S, Maezawa K, Kurosawa H. Activation of
263 fibrinolysis by reinfusion of unwashed salvaged blood after total knee
264 arthroplasty. *Transfus Apher Sci* 2010; 42: 33-37.
- 265 16. Collen D, Lijnen HR. Basic and clinical aspects of fibrinolysis and

- 266 thrombolysis. *Blood* 1991; 78: 3114-3124.
- 267 17. Hoylaerts M, Rijken DC, Lijnen HR, Collen D. Kinetics of the activation of
268 plasminogen by human tissue plasminogen activator. Role of fibrin. *J Biol*
269 *Chem* 1982; 257: 2912-2919.
- 270 18. Loscalzo J, Braunwald E. Tissue plasminogen activator. *N Engl J Med* 1988;
271 319: 925-931.
- 272 19. Fay WP, Eitzman DT, Shapiro AD, Madison EL, Ginsburg D. Platelets
273 inhibit fibrinolysis in vitro by both plasminogen activator inhibitor-1-dependent
274 and -independent mechanisms. *Blood* 1994; 83: 351-356.
- 275 20. Dellas C, Loskutoff DJ. Historical analysis of PAI-1 from its discovery to its
276 potential role in cell motility and disease. *Thromb Haemost* 2005; 93: 631-640.
- 277 21. Bruwer G, Limperger V, Kenet G, Klostermeier UC, Shneyder M,
278 Degenhardt F, et al. Impact of high risk thrombophilia status on recurrence
279 among children and adults with VTE: An observational multicenter cohort study.
280 *Blood Cells Mol Dis* 2016; 62: 24-31.
- 281 22. Di Minno MN, Ambrosino P, Ageno W, Rosendaal F, Di Minno G, Dentali F.
282 Natural anticoagulants deficiency and the risk of venous thromboembolism: a
283 meta-analysis of observational studies. *Thromb Res* 2015; 135: 923-932.
- 284 23. Kearon C. Natural history of venous thromboembolism. *Circulation* 2003;
285 107: I22-30.
- 286 24. Nilsson IM. Coagulation and fibrinolysis. *Scand J Gastroenterol Suppl* 1987;
287 137: 11-18.

- 288 25. Chmielewska J, Ranby M, Wiman B. Evidence for a rapid inhibitor to tissue
289 plasminogen activator in plasma. *Thromb Res* 1983; 31: 427-436.
- 290 26. Zhang Y, Wu TB, Daigle BJ, Jr., Cohen M, Petzold L. Identification of
291 disease states associated with coagulopathy in trauma. *BMC Med Inform Decis*
292 *Mak* 2016; 16: 124.
- 293 27. Hakki SI, Fareed J, Hoppensteadt DA, Abdullah H, Camblin J, Nasser AF,
294 et al. Plasma tissue factor pathway inhibitor levels as a marker for postoperative
295 bleeding after enoxaparin use in deep vein thrombosis prophylaxis in
296 orthopedics and general surgery. *Clin Appl Thromb Hemost* 2001; 7: 65-71.
- 297 28. Cosmi B. An update on the pharmaceutical management of thrombosis.
298 *Expert Opin Pharmacother* 2016: 1-16.
- 299 29. Gomez-Outes A, Terleira-Fernandez AI, Suarez-Gea ML, Vargas-Castrillon
300 E. Dabigatran, rivaroxaban, or apixaban versus enoxaparin for
301 thromboprophylaxis after total hip or knee replacement: systematic review,
302 meta-analysis, and indirect treatment comparisons. *BMJ* 2012; 344: e3675.

303 -----

304 **Table 1: Clinical characteristics in all TJA Patients.**

	TJA (n=119)	THA(n=48)	TKA(n=71)
Age (years)	57.8±12.0	63.5±7.9	49.3±12.2
Sex (male, %)	25.2	35.4	18.3
BMI (kg/m ²)	24.3±1.3	24.1±1.8	24.5±1.4
Hypertension (%)	35.3	22.9	43.7

Diabetes (%)	7.6	2.1	11.3
Cerebrovascular disease (%)	1.7	2.1	1.4
Coronary heart disease (%)	5.9	0	9.9
Use of glucocorticoids (%)	2.5	6.3	0
Joint replacement (%)	1.7	2.1	1.4

305 Values presented as mean±SD or n (%).

306 BMI, body mass index; TJA, total joint arthroplasty; THA, total hip
 307 arthroplasty; TKA, total knee arthroplasty; SD, standard deviation; TJA group
 308 includes both THA and TKA.

309 -----

310 **Table 2: Preoperative and Postoperative Values for coagulation and**
 311 **fibrinolytic biomarkers.**

	Pre	24h	72h	p-value (Pre vs 24h)	p-value (Pre vs 72h)
PC (µg/ml)	2.03±0.66	1.20±0.36	0.88±0.30	0.0176	0.00315
EPCR (µg/ml)	199.4±57.4	118.9±36.3	100.8±31.0	0.0446	0.00955
AT-III (µg/ml)	70.0±20.5	81.4±24.9	118.7±34.6	0.0129	0.00432
TFPI (µg/l)	16.68±8.92	20.14±7.94	26.37±7.91	0.0268	0.00677
tPA (ng/ml)	2.03±1.03	2.53±0.86	2.87±0.83	0.0409	0.00675
PAI-1 (ng/ml)	2.03±0.66	0.88±0.30	1.20±0.36	0.0201	0.00917

PC, Protein C; EPCR, Endothelial cell protein C receptor; AT-III, Antithrombin
 III; TFPI, Tissue factor pathway inhibitor; tPA, Tissue plasminogen activator;
 PAI-1, Plasminogen activator inhibitor 1.




Figure 1: Changes of the Protein C (PC) and Endothelial cell protein C receptor (EPCR) levels in patients compared to healthy controls in the pre- and postoperative period. Both the pre- and postsurgical plasma samples were significantly different in comparison to the control. The boxes represent the mean data ranges. The perpendicular lines represent standard deviation. Statistical differences were observed in the plasma levels of these biomarkers preoperatively and on postoperative 24h, 72h.

****denotes P < 0.05. *** denotes P < 0.01.**




Figure 2. Changes of the Antithrombin III (AT-III) and Tissue factor pathway inhibitor (TFPI) levels in patients compared to healthy controls in the pre- and postoperative period. Both the pre- and postsurgical plasma samples were significantly different in comparison to the control. The boxes represent the mean data ranges. The perpendicular lines represent standard deviation. Statistical differences were observed in the plasma levels of these biomarkers preoperatively and on postoperative 24h, 72h. **denotes $P < 0.05$. * denotes $P < 0.01$.**




Figure 3. Changes of the Tissue plasminogen activator (tPA) and Plasminogen activator inhibitor 1(PAI-1) levels in patients pre- and postoperative period compared to healthy controls. Both the pre- and postsurgical plasma samples were significantly different in comparison to the control. The boxes represent the mean data ranges. The perpendicular lines represent standard deviation. Statistical differences were observed in the plasma levels of these biomarkers preoperatively and on postoperative 24h, 72h. **denotes $P < 0.01$. * denotes $P < 0.001$.**