

1 **DOI : <https://doi.org/10.47391/JPMA.03-414>**

2

3 **Disclosing bad news of cancer diagnosis: patients' preference for**
4 **communication**

5

6 **Sabah Uddin Saqib, Khawaja Muhammad Inam Pal**

7 Department of Surgery, Aga Khan University Hospital, Karachi, Pakistan.

8 **Correspondence:** Sabah Uddin Saqib. **Email:** drsabahsaqib@hotmail.com

9

10 **Abstract**

11 The majority of relatives of cancer patients in Pakistan request their clinicians to
12 adopt a "do not tell approach" while counselling the patients regarding their
13 disease. The current study aimed to assess patients' understanding of their disease
14 and how they would prefer the physicians to deliver news about cancer diagnosis
15 and its management plan. This was a cross-sectional study in which both patients
16 and their immediate relatives were interviewed. The study enrolled 55 patients
17 with six different types of cancers. The study showed that 35 (65.5%) patients did
18 not know the stage of their illness at the time of diagnosis, while 40 (72.7%)
19 patients did not know the current stage of their disease. In 22 (40%) cases, the
20 patient's family knew the diagnosis ahead of the patient, and 19 (86.3%) families
21 asked the clinicians to hide the diagnosis from the patient. This study, which used
22 a scoring questionnaire, demonstrates that specialist oncologists for breaking the
23 bad news, family counselling, helping patients to figure out how to inform others,
24 giving the news directly to the patient and the effects of cancer on daily life are
25 preferred areas to communicate with cancer patients.

26 **Keywords:** Breaking news, Cancer, patients' Preference, Communication

27

28

29 **Introduction**

30 Bad news in medical literature is usually defined as any news which negatively
31 changes or alters the views of a patient, regarding his or her life or future.¹ In the
32 last few decades, there has been a lot of emphasis on patient autonomy,
33 empowerment, and the patient's direct involvement in decision-making regarding
34 the management of their illnesses, especially if they are suffering from cancer.²
35 As a result of this, now in most of the developed countries, all information related
36 to health and disease is disclosed first to the patient.³ However, it is still widely
37 seen in many cultures that breaking bad news of cancer is a disagreeable
38 experience for patients, and most of the time it causes many patients to lose hope.
39 In many instances, the patient's family is conveyed the bad news, while the
40 patient is not present at the time of disclosure of any unfavourable news. This
41 happens mostly because the family wishes or at times the patients' relatives put
42 undue pressure on the clinicians to not to tell the bad news to the patients directly,
43 whether the bad news is about diagnosis, the need for an intense treatment, or
44 poor prognosis. Unfortunately, in these cultures physicians are often seen to
45 follow the family's wishes.¹

46 Cancer is currently one of the leading cause of death and prolonged illness; it
47 resulted in 14 million new cases and 8.2 million deaths, worldwide in 2012.⁴ On
48 an average, with each passing minute, 16 patients lose their lives in their fight
49 against cancer, while 26 new individuals are informed that they have some form
50 of cancer.⁵ Diagnosis of cancer is always perceived as unfavourable news as not
51 only the patients but the whole family is affected. Debilitating and disfiguring
52 treatment, pain, loss of function, costly treatment, indefinite follow-ups,
53 recurrences, and death are the main issues of discussion. Although in every field
54 of medicine, clinicians often have to communicate adverse medical information
55 to the patients, it is particularly common in oncology setting where unfavourable
56 news of life-threatening conditions, unwanted treatment, and often disease
57 recurrence has to be communicated to the patients and their families. This

58 communication often ends up on a very difficult question to an oncologist: how
59 much time do we have? Even the best specialist finds this question very hard to
60 answer.

61 Pakistan is a developing, low to middle-income country where modern treatment
62 options for treating cancer are either not readily available or they are very costly.⁶
63 Secondly, low literacy rates and myths attached to the disease often make a
64 diagnosis and available treatment options very difficult to understand not only for
65 the patients but also for their families.⁷ There are guidelines available which
66 illustrate how unfavourable news should be communicated and unwanted
67 interviews should be conducted. But in countries like Pakistan, where cancer is
68 still considered the biggest taboo, reactions while admitting bad news of cancer
69 can be very unpredictable both for the patient and their families.

70 Many clinicians, whether they belong to surgical, medical or radiation oncology,
71 firmly believe that the patient has every right to know about his or her illness and
72 only this knowledge can help them to sail through the various periods of the
73 difficult journey. Few studies are available which have assessed the patients'
74 reactions to how bad news of cancer was conveyed to them or have analysed the
75 concurrent relationship between physicians and patients, in terms of how both
76 rank the way bad news was delivered.⁸⁻¹⁰ Since delivering bad news of cancer is
77 a two-way communication between the physician and the patient, and as the
78 patient is the one whose life is directly affected, it is very important to consider
79 the patient's understanding, fears, and preferences for communication while
80 conducting such interviews in any clinical setting. Thus, our study aimed to assess
81 the patients' preferences regarding how they should be informed not only about
82 the diagnosis of cancer but also about various management options available.

83

84 **Patients and methods**

85 This was a cross-sectional study, conducted from October 1, 2016 to December
86 31, 2016 at the outpatient department rooms of Medical and Surgical Oncology,

87 Aga Khan University Hospital, Karachi. All participants in this study were aged
88 18 or above and they were interviewed during their follow-up appointments in
89 the medical or surgical clinics. We used convenient sampling technique and only
90 those patients were included, who were diagnosed with cancer at least one month
91 earlier to allow them time to adjust to the news of diagnosis and to reflect on their
92 experience. The study was briefly described, and they were asked to participate.
93 The patients who consented to participate were enrolled. All individuals were
94 interviewed in a separate room in their respective clinics and were given a
95 researchers' self-designed simple questionnaire covering different aspects of
96 communication; they were asked to rate the importance of different
97 communicating methods while counselling on cancer in terms of score — score
98 1 being not at all effective method and score 5 highly important skills. This rating
99 scale questionnaire was available in both Urdu and English languages. Fourteen
100 questionnaire items were selected to assess patients' preference for
101 communication. Score 4 or higher was taken as the best way of communication.
102 Score less than 3 was regarded as a bad way of communication, while score 3-4
103 was regarded as an intermediate skill of communication, which can be used
104 according to the situation. The confidentiality of all patients was maintained. The
105 primary investigator also asked some questions to assess the patients'
106 understanding regarding their disease; an immediate family member was also
107 questioned regarding their understanding.

108

109 **Results**

110 Seventy patients were approached to participate in this study. Twelve patients
111 declined to participate as they were already exhausted due to long waiting times
112 in the clinic waiting area. Three patients did not participate in the study because
113 they did not want to recall the bad experience, which they had in the past when
114 they were first told about the disease. Hence, in total 55 patients gave informed

115 consent before participating in the study. The mean interval (SD) between the
116 diagnosis of their disease and interview was 4.7 months(± 3.7)

117 In our study, out of 55 patients, 25 (45.5%) were men and 30(54.5%) were
118 women. The mean age of the interviewed population was 53 ± 13 years. Out of the
119 55 patients, 18(32.7%) were diagnosed with gastrointestinal cancer, 6(10.9%)
120 with urinary tract cancer, 4(7.3%) with lung cancer, 9(16.4%) with
121 gynaecological cancer, 8(14.5%) with breast cancer, and 10(18.2%) with head
122 and neck cancer. On average, three doctors per patient were visited before
123 establishing the actual diagnosis of cancer. Moreover, the mean time from the
124 onset of symptoms to the diagnosis was 4.85 ± 2.5 months. (Table 1)

125 When the patients were asked about their knowledge related to their disease
126 45(81.8 %) patients could tell the name of the organ which had developed cancer.
127 Only 19(34.5%) of the patients knew the stage of cancer at the time of diagnosis,
128 while only 15(27.3 %) patients were aware of their current stage of cancer (Table
129 2). Around 40 out of 55(73%) patients never thought that their symptoms might
130 be because of cancer. When patients were first told regarding their disease,
131 20(36%) patients completely rejected this diagnosis, 13(24%) started thinking
132 about living, 6(11%) thought about family, 6(11%) thought they should have
133 been told much earlier, while only 10(18%) patients were optimistic to take
134 treatment.

135 In our study, the approach that only cancer specialist should break the news of
136 cancer (mean score 4.2 ± 0.7 , news should be told directly to the patient first (e
137 4 ± 1), clinicians should offer some hope to the patients (mean score 4 ± 0.9), a
138 patient's family should be informed simultaneously (mean score 4.2 ± 0.8),
139 educating patients about how cancer affects daily living and how patients can talk
140 to others regarding their disease (mean score 4 ± 1), were regarded as the best skill
141 of communication and every clinician should approach cancer patients in this
142 manner. Our survey results completely rejected the approach of warning the
143 patients first before breaking the news (mean score 2.5 ± 1.3), waiting to disclose

144 cancer news till all staging workup is available (mean score 2.7 ± 1.2), blunt
145 behaviour of doctor (mean score 1.7 ± 0.9) and a crowded room with some
146 teaching for students, while communicating with cancer patients (mean score 1.6
147 ± 0.8). (Table 3). We also interviewed the patient's next of kin and the results
148 showed that in 20 out of 55 cases (36%) patient's family knew diagnosis ahead
149 of patient and 19 (34.5%) families requested their clinicians to hide the news of
150 cancer from the patients. Out of these 19 families 8(42%) thought that it will add
151 anxiety, another 8(42%) families thought that cancer news is itself fearful news,
152 while 3(16%) families believed that the patient was not ready for this news (Table
153 4).

154

155 **Discussion**

156 The results of our study showed that in our population a significant amount of
157 time is spent on establishing the diagnosis of cancer. Besides, lack of screening,
158 fear, and poor communication are the other factors of delay, which result in a
159 change of multiple doctors. This study showed the preferences of patients for
160 communication, based on their experience, when they were first told about
161 cancer. The results of our survey demonstrate that there is certain information
162 which the cancer patients want to be given directly to them at a certain point and
163 also some recommendations for clinicians for improving communication with
164 cancer patients. A study done by Sardell and Trierweiler highlighted the
165 importance of hopefulness in the course of cancer treatment.¹¹ The patients in our
166 study showed their personal preferences regarding how they would like the news
167 conveyed to them, and their ratings showed that content of the message, setting
168 where the news is communicated and core knowledge of a clinician is all-
169 important, and their rating for communication of cancer news is almost similar to
170 the study by Parker et al which highlighted the importance of these fundamentals
171 of good communication.¹² Gebhardt et al¹³, in their study, concluded that
172 physicians should disclose bad news in a patient-centered way to reduce

173 mismatch with patients' preferences and help them in reducing the anxiety and
174 stress associated with unfavourable news of cancer. Our study also disregards the
175 preference of family members to hide the news of cancer from the patients and
176 concluded that communication regarding disease and disclosure of any news
177 should be patient-centred.

178 Our study had some limitations, i.e. it was done in a tertiary care hospital with
179 advanced oncology units and skilled professionals, hence it does not take into
180 account those patients who are seen for cancer in small hospitals of our country.
181 Although the study had a small sample size, it had a diverse group of patients, not
182 only from the disease point of view but also from their socio-economic
183 background, which gives a good foundation to conduct a study with a large
184 sample size prospectively.

185

186 **Conclusion**

187 Our study concluded that the patients have specific preferences for
188 communication at the time of disclosure of bad news of cancer and the clinician
189 must take into account such considerations. As clinicians, we should not use those
190 communication tools which were highly disregarded by the patients based on
191 their personal experiences.

192

193 **Acknowledgments:** Medical, surgical, radiation oncologists of Aga Khan
194 University Hospital, Karachi, for giving me access to interview their patients
195 during their busy clinics. All nurses of the oncology department for their co-
196 operation in this study.

197 **Disclaimer:** None to declare.

198 **Conflict of Interest:** None to declare.

199 **Funding Sources:** No funding was involved in this study.

200 **Ethical approval:** Ethical review committee of AKUH approved this study and
201 approval no is 4403-Sur- ERC-2016

References

- 202 1. Monden KR, Gentry L, Cox TR. Delivering bad news to patients. Proc
203 (Bayl Univ Med Cent). 2016;29:101-2.
- 204 2. Murgic L, Hebert PC, Sovic S, Pavlekovic G. Paternalism and autonomy:
205 views of patients and providers in a transitional (post-communist) country. BMC
206 Med Ethics. 2015;16:65.
- 207 3. Nass SJ, Levit LA, Gostin LO. The value and importance of health
208 information privacy. Beyond the HIPAA Privacy Rule: Enhancing Privacy,
209 Improving Health Through Research: National Academies Press (US); 2009.
- 210 4. Stewart BW, Bray F, Forman D, Ohgaki H, Straif K, Ullrich A, et al.
211 Cancer prevention as part of precision medicine: 'plenty to be done'.
212 Carcinogenesis. 2016;37:2-9.
- 213 5. Ferlay J, Soerjomataram I, Dikshit R, Eser S, Mathers C, Rebelo M, et al.
214 Cancer incidence and mortality worldwide: sources, methods and major patterns
215 in GLOBOCAN 2012. Int J Cancer. 2015;136:E359-86.
- 216 6. Zafar SN, McQueen KA. Surgery, public health, and Pakistan. World J
217 Surg. 2011;35:2625-34.
- 218 7. Carlsson ME, Strang PM. Facts, misconceptions, and myths about cancer.
219 Gynecologic oncology. 1997;65:46-53.
- 220 8. Maguire P. Improving communication with cancer patients. European
221 Journal of Cancer. 1999;35:2058-65.
- 222 9. Ong LM, Visser MR, Lammes FB, De Haes JC. Doctor-patient
223 communication and cancer patients' quality of life and satisfaction. Patient
224 education and counseling. 2000;41:145-56.
- 225 10. Peterson EB, Ostroff JS, DuHamel KN, D'Agostino TA, Hernandez M,
226 Canzona MR, et al. Impact of provider-patient communication on cancer
227 screening adherence: a systematic review. Preventive medicine. 2016;93:96-105.
- 228 11. Sardell AN, Trierweiler SJ. Disclosing the cancer diagnosis. Procedures
229 that influence patient hopefulness. Cancer. 1993;72:3355-65.
- 230

231 12. Parker PA, Baile WF, de Moor C, Lenzi R, Kudelka AP, Cohen L.
 232 Breaking bad news about cancer: patients' preferences for communication. J Clin
 233 Oncol. 2001;19:2049-56.

234 13. Gebhardt C, Gorba C, Oechsle K, Vehling S, Koch U, Mehnert A.
 235 [Breaking Bad News to Cancer Patients: Content, Communication Preferences
 236 and Psychological Distress]. Psychother Psychosom Med Psychol. 2017;67:312-
 237 21.

238

239 -----

240

241 **Table 1: Demographics and medical characteristics of patients**

Gender	Number (%)
Male	25(45.5)
Female	30(54.5)
Mean Age years	53 ± 13
Number of doctors on average visited before diagnosis	3 per patient
Meantime since diagnosis (months)	4.7 ±3.7
Meantime elapsed from the onset of symptoms until the establishment of the diagnosis (months)	4.85 ±2.5
Type of cancer	Number (%)
Gastrointestinal cancers	18 (32.7)
Urinary tract cancers	6 (10.9)
Lung cancers	4 (7.3)
Gynecological cancers	9 (16.4)
Breast cancers	8 (14.5)
Head and neck cancers	10 (18.2)

242

243 -----

244

245

246

247

248

249 **Table 2: Level of understanding of patients regarding their disease**

	Number	(%)
Patients knowing organ name from which cancer originated	45	(81.8)
Patients knowing stage at diagnosis	19	(34.5)
Patients knowing the current stage of cancer	15	(27.3)

250

251 -----

252

253 **Table 3: Highest and lowest ratings of patients' preferences regarding**
254 **different aspects of cancer news delivery**

Item	Mean	SD ±
Highest rating questions		
The doctor who breaks news should be a specialist of that particular cancer	4.27	0.7
The doctor tells news directly to the patient	4	1
The doctor tells, how cancer can affect my daily living	4.2	0.7
Having a doctor offering me hope	4	0.9
The doctor informs my family about my diagnosis	4.2	0.8
Doctor helps me how to tell others about my diagnosis	4	1
Intermediate rating questions		
The doctor tells all treatment options on the first encounter	3.5	1.5
The doctor tells me, my 5 years survival chance	3	1.4
The doctor should tell diagnosis as soon as biopsy report is available	3	1.6
Doctor asks me to bring close relative before breaking news	3.3	1.2
Lowest rating questions		
Doctor warns me that he has unfavourable news	2.5	1.3
News of cancer should wait till staging workup is available	2.7	2
The doctor is very blunt and not showing empathy	1.7	0.9
Doctor not making eye contact and teaching his or her students	1.6	0.8

255

256 -----

257

258 **Table 4: Response of family members after knowing the diagnosis**

The response of family members	Number (%)
Number of families knew diagnosis before the patient	20(36%)
Number of families requested the doctor to hide the news from the patient	19(34.5%)
The reason behind hiding news from the patient	
News will cause anxiety to the patient	8(42%)
Cancer news is a fearful news	8(42%)
The patient is not mentally prepared	3(16%)

259

Provisionally Accepted for Publication