

1 **DOI: <https://doi.org/10.47391/JPMA.04-593>**

2
3 **Correlation of serum calcium with severity of acute ischaemic stroke**

4
5 **Ainan Arshad¹, Saad Bin Zafar Mahmood², Ahmed Ayaz³, Mehssa Hashim⁴, Eisha**
6 **Hanan⁵**

7 **1,2** Department of Medicine, Aga Khan University Hospital, Karachi, Pakistan; **3** Department of
8 Paediatrics, Aga Khan University Hospital, Karachi, Pakistan; **4,5** MBBS 3rd Year Student, Medical
9 college, Aga Khan University Hospital, Karachi, Pakistan.

10 **Correspondence:** Ainan Arshad. **Email:** ainan_arshad@hotmail.com

11
12 **Abstract**

13 Stroke results in the death of around 6.5 million people annually with a majority of these
14 occurring in developing countries. Serum calcium has been hypothesised to play a
15 significant role in causing ischaemic stroke. This retrospective observational study was
16 conducted to determine the correlation, if any, between serum calcium and the severity of
17 acute ischaemic stroke in our population. Two hundred and seventy-nine patients admitted
18 with acute ischaemic stroke were enrolled in the study. Of the 279 patients 162 (58%) were
19 male and mean age was 62.4 ± 3.8 years. Characteristics of stroke patients were compared
20 with stroke severity. Mean albumin corrected serum calcium and Scandinavian stroke
21 severity score was $9.1 (\pm 5.6)$ and $33.67 (\pm 15.2)$, respectively. Hypertension and mean
22 GCS on admission were significantly associated with increased stroke severity score.
23 However, no correlation was observed between serum calcium and severity of acute
24 ischaemic stroke.

25 **Keywords:** Stroke, Serum calcium, Scandinavian stroke severity scale, cerebral ischaemia

27 **Introduction**

28 Stroke is categorised as the number one cause of disability and the third most common
29 cause of mortality worldwide. Literature reports that an estimated 6.5 million people lose
30 their lives due to stroke every year, out of which 75% deaths occur in developing countries.¹

31 The majority of strokes in Pakistan are a result of cerebral ischaemia, with the most
32 common type being lacunar stroke.² TOAST (Trial of Org 10172 in Acute Stroke
33 Treatment) classification of stroke has classified the causes of ischaemic stroke into: large
34 artery atherosclerosis, cardio-embolism, small vessel disease, other determined aetiologies
35 and undetermined causes.³ Multiple variables have an impact on the prognosis and outcome
36 of stroke, such as age, comorbidities, cause of stroke, location and severity of stroke, level
37 of consciousness and blood glucose.³

38 Calcium is known to play a significant role in the pathogenesis of ischaemic stroke, since
39 cerebral ischaemia causes an increase in intracellular calcium resulting in activation of
40 cytotoxic enzymes leading to cell death.⁴ The present study was conducted to determine the
41 correlation, if any, between serum calcium and severity of acute ischaemic stroke, and to
42 establish the local perspective as there is a paucity of local data even though we are
43 burdened with a large number of people presenting with acute ischaemic stroke.

44

45 **Patients/Methods and Results**

46 A retrospective observational study was conducted at the Department of Medicine, Aga
47 Khan University Hospital, Karachi, from January 2018 to December 2018, after taking
48 ethical approval from the Ethical Review Committee (2018-0267-153). All patients
49 presenting acutely within 24 hours of suffering stroke with CT scan at the time of admission
50 showing a hypodense area, plus any two or more of the following symptoms inability to
51 move one or more limb, aphasia, imbalance and GCS <15, were labelled as acute ischaemic
52 stroke. Severity of the stroke was measured according to the Scandinavian stroke scale
53 (SSS). Patients with a history of mania, bipolar effective disorder or post-traumatic stress,

54 history of thyroid and parathyroid disorders, history of blood transfusion within one week,
55 history of CNS disease (e.g. head trauma, multiple sclerosis), and with history of congestive
56 heart failure, chronic renal failure, chronic obstructive pulmonary disease and chronic liver
57 disease, were excluded. A score of less than 43 was classified as mild stroke, 26-43 as
58 moderate, and less than 26 as severe stroke.⁵

59 We compared patients' characteristics as mild, moderate, and severe stroke by using
60 Student t-test or Wilcoxon rank-sum test for continuous variables and chi square test or
61 Fisher exact test for categorical variables based on cell counts. Spearman rank correlation
62 test was performed to investigate the relationship between calcium quartiles and stroke
63 severity and p-value of ≤ 0.05 was taken as significant.

64 A total of 730 patients were reviewed out of which 279 fulfilled the inclusion criteria. Of
65 the 279 patients, 162 (58%) were male, while 117 (42%) were female. The mean age of the
66 sample population was 62.4 ± 13.8 with age range between 19 and 94 years. A total of 211
67 (75.6%) patients were hypertensive with a mean systolic BP of 150.1 ± 36.0 mmHg and
68 diastolic BP of 83.2 ± 18.6 mm Hg. Mean GCS on admission to the hospital was found to
69 be 12.7 ± 3.2 . Mean serum calcium level at admission (adjusted for albumin) and mean
70 Scandinavian stroke scale severity score were 9.1 ± 5.6 and 33.67 ± 15.2 , respectively. The
71 rest of the baseline characteristics have been summarised in Table 1.

72 The sample patient population was almost equally divided into three categories of stroke
73 severity scale (Mild = 90, Moderate = 96, and Severe = 93). Mean serum calcium level at
74 the time of admission was not significantly related to stroke severity. However, baseline
75 comorbid and mean GCS on admission were both significantly related to stroke severity (p
76 value < 0.05). Table 2 shows the characteristics of stroke patients according to stroke
77 severity score.

78

79

80

81 **Discussion**

82 Stroke is one of the foremost causes of mortality and morbidity throughout the world,
83 whether it is in the developing or developed country.¹ Calcium ions are known to be
84 involved in the pathophysiology of cerebral ischaemia as hypoxia/ischaemia results in
85 transfer of calcium ions from extracellular to intracellular cerebral tissue spaces.⁴ Our
86 study, however, showed no significant correlation between serum calcium and acute
87 ischaemic stroke severity.

88 Review of literature, both global as well as regional has shown conflicting results. Some
89 studies have found serum calcium to be low in transient ischaemic attack and further low
90 calcium has been hypothesised to occur in ischaemic cerebral infarction.⁶ Some meta-
91 analyses have even reported an inverse association between dietary calcium intake and risk
92 of ischaemic stroke.⁷ However, some studies have reported that there is no significant
93 association between serum calcium and stroke severity or serum calcium and risk of
94 ischaemic stroke.⁸⁻⁹ One study has even suggested that higher levels of serum calcium are
95 associated with more severe artery stenosis lesions.¹⁰ These arguments have led us to
96 believe that serum calcium is not a reliable marker for accurate prediction of severity of
97 stroke or its prevalence.

98 The study is limited by its design (retrospective study) and the timing of calcium samples
99 could also not be accurately ascertained. However, it is important to note that all patients
100 had a serum calcium sent at admission as part of initial baseline workup, from the
101 emergency. Furthermore, since a majority of the patients presenting to our tertiary hospital
102 have either been previously admitted elsewhere or are from far flung areas, true serum
103 calcium values can neither be found nor can it be judged, if the value of serum calcium will
104 be classified as early or late even if the study was done prospectively. Another problem
105 with retrospective data collection was the quality of the data in the file and variability in
106 the examination done by the physicians. Nonetheless, this study has offered new local
107 information to clinicians which can influence clinical practice. Previous studies on this

108 topic have used the modified NIHSS scale. Our study is unique as it uses the Scandinavian
109 stroke severity scale, which has been found to be equal to the NIHSS and is easy to apply
110 in the acute clinical setting.⁵

111

112 **Conclusion**

113 In conclusion, our results did not show any correlation between serum calcium levels at the
114 time of admission and ischaemic stroke severity. However, further studies in South Asian
115 population are warranted to properly quantify the correlation between serum calcium and
116 stroke severity and to assess the importance of serum calcium as a prognostic marker of
117 stroke outcome.

118

119 **Disclaimer:** None

120 **Conflict of interest:** The authors declare they have no conflict of interest

121 **Funding disclosure:** None

122

123 **References**

- 124 1. Venketasubramanian N, Yoon BW, Pandian J, Navarro JC. Stroke Epidemiology in
125 South, East, and South-East Asia: A Review. *Journal of stroke*. 2017;19:286-94.
- 126 2. Zafar F, Tariq W, Shoaib RF, Shah A, Siddique M, Zaki A, et al. Frequency of
127 Ischemic Stroke Subtypes Based on Toast Classification at a Tertiary Care Centre in
128 Pakistan. *Asian journal of neurosurgery*. 2018;13:984-9.
- 129 3. Radu RA, Terecoasa EO, Bajenaru OA, Tiu C. Etiologic classification of ischemic
130 stroke: Where do we stand? *Clinical neurology and neurosurgery*. 2017;159:93-106.
- 131 4. Chung JW, Ryu WS, Kim BJ, Yoon BW. Elevated calcium after acute ischemic
132 stroke: association with a poor short-term outcome and long-term mortality. *Journal*
133 *of stroke*. 2015;17:54-9.

- 134 5. Labberton AS, Rønning OM, Thommessen B, Barra M. Changes in survival and
 135 characteristics among older stroke unit patients-1994 versus 2012. *Brain and*
 136 *behavior*. 2019;9:e01175.
- 137 6. Ishfaq M, Ullah F, Akbar S, Rahim F, Afridi AK. Correlation of serum calcium with
 138 severity of acute ischaemic stroke. *J Pak Med Assoc*.2017;67:20-3.
- 139 7. Dibaba DT, Xun P, Fly AD, Bidulescu A, Tsinovoi CL, Judd SE, et al. Calcium
 140 Intake and Serum Calcium Level in Relation to the Risk of Ischemic Stroke: Findings
 141 from the REGARDS Study. *Journal of stroke*. 2019;21:312-23.
- 142 8. Guven H, Cilliler AE, Koker C, Sarikaya SA, Comoglu SS. Association of serum
 143 calcium levels with clinical severity of acute ischemic stroke. *Acta neurologica*
 144 *Belgica*. 2011;111:45-9.
- 145 9. Larsson SC, Traylor M, Burgess S, Boncoraglio GB, Jern C, Michaëlsson K, et al.
 146 Serum magnesium and calcium levels in relation to ischemic stroke. *Neurology*.
 147 2019;92:e944.
- 148 10. Wu J, Xie J, Zhao Y, Gong L, Liu X, Liu W. Serum Calcium is Related to the Degree
 149 of Artery Stenosis in Acute Ischemic Stroke. *Cellular physiology and biochemistry:*
 150 *International Journal of Experimental Cellular Physiology, Biochemistry, and*
 151 *Pharmacology*. 2018;46:1189-97.

152
 153 -----
 154
 155 **Table 1: Characteristics of stroke patients (N=279)**

Characteristics	Overall (N = 279)
Mean age (years)	62.4±13.8
Gender	
Male	162 (58%)
Female	117 (42%)
Comorbid conditions	

Diabetes	152 (54.5%)
Hypertension	211 (75.6%)
Smoking	33 (11.8%)
Hospitalisation factors	
Mean duration of symptoms (hours)	32.9 ± 36.1
Mean systolic BP on admission(mm Hg)	150.1 ± 36.0
Mean diastolic BP on admission(mm Hg)	83.2 ± 18.6
Mean GCS on admission	12.7 ± 3.2
Mean Haemoglobin on admission (Gm/dl)	13.8 ± 10.8
Mean Total Cholesterol (mg/dl)	146.6 ± 62.7
Mean Triglycerides (mg/dl)	125.2 ± 79.3
Mean HDL (mg/dl)	35.65 ± 16.7
Mean LDL (mg/dl)	90.27 ± 45.8
Mean Serum Calcium (mg/dl)	9.1 ± 5.6
Mean Stroke severity score	33.67 ± 15.2
Stroke Severity	
Mild	90 (32.3%)
Moderate	96 (34.4%)
Severe	93 (33.3%)

156 BP: Blood pressure, GCS: Glasgow coma score, HDL: High density lipoprotein, LDL: Low
 157 density lipoprotein
 158

159

160

161

Table 2: Characteristics of stroke patients according to stroke severity score (N=279)

Characteristics	Overall (N = 279)	Mild (N = 90)	Moderate (N = 96)	Severe (N = 93)	P- value
Mean age (years)	62.4±13.8	59.9±14.3	64.9±12.6	62.19±14.3	0.547
Gender					0.268
Male	162(58%)	58	55	49	
Female	117(42%)	32	41	44	
Comorbid conditions					
Diabetes	152(54.5%)	41	55	56	0.109
Hypertension	211(75.6%)	58	79	74	0.010

Smoking	33(11.8%)	9	12	12	0.805
Hospitalisation factors					
Mean duration of symptoms (hours)	32.9 ±36.1	33.24± 35.9	32.47± 36.7	33.12± 36.1	0.595
Mean systolic BP on admission(mm Hg)	150.1±36.0	152.57 ± 50.0	154.16±27.6	143.37±25.4	0.181
Mean diastolic BP on admission(mm Hg)	83.2± 18.6	83.3± 17.3	84.01± 18.1	82.42± 20.5	0.637
Mean GCS on admission	12.7± 3.2	14.5± 1.8	13.6± 2.0	10.0± 3.5	<0.001
Mean Haemoglobin on admission	13.8±10.8	13.0± 2.4	13.8±12.4	14.5±13.7	<0.001
Mean Total Cholesterol(mg/dl)	146.6±62.7	152.6± 56.3	151.0± 57.5	136.1±72.4	0.657
Mean Triglycerides(mg/dl)	125.2±79.3	137.9± 85.9	122.4± 66.2	115.8±84.4	0.614
Mean HDL(mg/dl)	35.65±16.7	36.2±14.5	37.1±19.1	33.6±19.1	0.463
Mean LDL(mg/dl)	90.27±45.8	94.2± 40.7	93.5±44.8	83.1±50.8	0.766
Mean Serum Calcium	9.1± 5.6	8.8± 0.5	9.7± 9.4	8.8±1.0	0.756

162 BP: Blood pressure, GCS: Glasgow coma score, HDL: High density lipoprotein, LDL: Low
 163 density lipoprotein
 164